

University of Groningen

Catalytic enantioselective synthesis of (-)-prostaglandin E-1 methyl ester based on a tandem 1,4-addition-aldol reaction

Arnold, L.A.; Naasz, R.; Minnaard, A.J.; Feringa, B.L.

Published in:
Journal of Organic Chemistry

DOI:
[10.1021/jo025987x](https://doi.org/10.1021/jo025987x)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2002

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Arnold, L. A., Naasz, R., Minnaard, A. J., & Feringa, B. L. (2002). Catalytic enantioselective synthesis of (-)-prostaglandin E-1 methyl ester based on a tandem 1,4-addition-aldol reaction. *Journal of Organic Chemistry*, 67(21), 7244-7254. <https://doi.org/10.1021/jo025987x>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Catalytic Enantioselective Synthesis of (-) Prostaglandin E₁ Methyl Ester Based on a Tandem 1,4-Addition-Aldol Reaction

Leggy A. Arnold, Robert Naasz, Adriaan J. Minnaard and Ben L. Feringa*

Department of Organic and Molecular Inorganic Chemistry, Stratingh Institute, University of Groningen, Nijenborgh 4, 9747 AG Groningen, The Netherlands
feringa@chem.rug.nl

Supporting information

Contents:

Characterization of compounds **7a** – **7c**, **7e**, **8**, **8a**, **8b**, **9**, **11a**, **11b**, **13a** – **15a**, **17a** – **21a**, **11c**, **13c** – **15c**, **17c**, **13d**, **20d**.

NOESY-NMR of compounds **11a**, **11b**, **28**

CD-spectra of compound **28**

Copy of ¹³C-NMR-spectra of compounds: **7a** - **7e**, **8** - **10**, **11a**, **11b**, **13a** – **21a**, **11c**, **13c** – **17c**, **13d**, **20d**, **21d**, **22** – **38**

General procedure for the acetalization of 2-cyclopenten-1,3-dione (7a-7e). To a cooled solution (0°C) of cyclopenten-3,5-dione (1.92 g, 20 mmol) and BF₃·Et₂O (2.52 ml, 20 mmol) in chloroform (50 ml) was added the alcohol (40 mmol, 20 mmol for diol). After stirring for 3 h the reaction mixture was poured into NH₄Cl (aq) and extracted three times with 25 ml diethyl ether. The combined organic layers were dried over MgSO₄, filtered and concentrated *in vacuo*. Column chromatography (SiO₂ ether/pentane) gave the corresponding acetals.

4,4-Dimethoxy-2-cyclopenten-1-one (7a). Purification by column chromatography (SiO₂ hexane/diethyl ether, 1:3, R_f = 0.37) gave 710 mg (26%) of **7a** as a colorless liquid. ¹H-NMR (300 MHz) δ = 7.52 (d, *J* = 6.0 Hz, 1H), 6.17 (d, *J* = 5.7 Hz, 1H), 3.28 (s, 6H),

2.49 (s, 2H); ^{13}C -NMR (200 Hz) δ = 203.6, 156.7, 135.1, 106.4, 50.2, 44.9, Ms (CI) for $\text{C}_7\text{H}_{10}\text{O}_3$: m/z = 142 (M^+), 160 ($\text{M} + \text{NH}_4^+$).

8,8-Dimethyl-6,10-dioxaspiro[4.5]dec-3-en-2-one (7b). Purification by column chromatography (SiO_2 hexane/diethyl ether, 1:1, R_f = 0.32) gave 1.03 g (28%) of **7b** as a white solid. Mp: 66-67°C; ^1H -NMR (300 MHz) δ = 7.54 (d, J = 6.0 Hz, 1H), 6.21 (d, J = 6.0 Hz, 1H), 3.60 (d, J = 16.0 Hz, 2H), 3.50 (d, J = 16.0 Hz, 2H), 2.60 (s, 2H), 1.09 (s, 3H), 0.89 (s, 3H); ^{13}C -NMR (200 MHz) δ = 204.1, 157.1, 135.3, 103.9, 72.9, 44.0, 29.9, 22.3, 22.2; HRMS calc. for $\text{C}_{10}\text{H}_{14}\text{O}_3$ 182.094, found 182.094; Anal.Calc. for $\text{C}_{10}\text{H}_{14}\text{O}_3$: C, 65.90; H, 7.74 found C, 65.98; H, 7.79.

1,4-Dioxaspiro[4.4]non-8-en-7-one (7c). Purification by column chromatography (SiO_2 hexane/diethyl ether, 1:2, R_f = 0.32) gave 907 mg (22%) of **7c** as a colorless liquid. ^1H -NMR (300 MHz) δ = 7.20 (d, J = 6.0 Hz, 1H), 6.17 (d, J = 6.0 Hz, 1H), 4.02 (m, 4H), 2.57 (s, 2H); ^{13}C -NMR (200 MHz) δ = 204.2, 156.5, 135.6, 111.9, 65.4, 45.4; Ms (CI) for $\text{C}_7\text{H}_8\text{O}_3$: m/z = 140 (M^+), 158 ($\text{M} + \text{NH}_4^+$).

2,2,3,3-Tetramethyl-1,4-dioxaspiro[4.4]non-8-en-7-one (7e). Purification by column chromatography (SiO_2 hexane/diethyl ether, 1:4, R_f = 0.38) gave 1.143 g (29%) of **7e** as a white solid. Mp: 59-60°C; ^1H -NMR: (300 MHz) δ = 7.17 (d, J = 6.0 Hz, 1H), 6.14 (d, J = 6.0 Hz, 1H), 2.60, (s, 2H) 1.25 (s, 6H), 1.23 (s, 6H); ^{13}C -NMR: (200 MHz) δ = 204.9, 159.6, 135.1, 109.0, 83.8, 48.4, 23.4, 23.2; Ms (CI) for $\text{C}_{11}\text{H}_{16}\text{O}_3$: m/z = 197 (M^+) 214 ($\text{M} + \text{NH}_4^+$).

General procedure for the 1,4-addition (8-10). A solution of $\text{Cu}(\text{OTf})_2$ (3.6 mg, 0.01 mmol) and phosphoramidite (0.02 mmol) in toluene (7 ml) was stirred under a nitrogen atmosphere at ambient temperature for 1 h. The cyclopenten-3,5-dione monoacetal (0.5

mmol) was added and after cooling the reaction mixture to -45 °C, the diorganozinc compound (0.6 ml of a 1 M solution in toluene) was added and stirring at -45°C was continued for 18 h. The conversion was determined by TLC. After complete conversion, the reaction mixture was poured in 25 ml NH₄Cl (aq) and the organic layer was separated, and the aqueous layer was extracted two times with diethyl ether. The combined organic layers were dried over MgSO₄, filtered and concentrated *in vacuo*.

(4*R*)-4-Ethyl-8,8-dimethyl-6,10-dioxaspiro[4.5]decane-2-one (8). Purification by column chromatography (SiO₂ pentane/diethyl ether, 7:1, R_f = 0.6) gave 33 mg (31%) of **8** as a colorless liquid. ¹H-NMR (300 MHz) δ = 3.45 (s, 4H), 2.96 (d, *J* = 18.0 Hz, 1H), 2.49 (dd, *J* = 17.7 Hz, *J* = 17.4 Hz, 1H), 2.35 (d, *J* = 18.0 Hz, 1H), 2.22 (m, 1H), 2.07 (dd, *J* = 17.7 Hz, *J* = 17.2 Hz, 1H), 1.89 (m, 1H), 1.39 (m, 1H), 1.16 (s, 3H), 0.94 (t, *J* = 7.8 Hz, 3H), 0.76 (s, 3H); ¹³C-NMR (200 MHz) δ = 213.5, 104.1, 72.8, 71.3, 47.7, 44.3, 42.7, 30.2, 22.6, 22.0, 20.6, 12.5; HRMS calc. for C₁₂H₂₀O₃ 212.141 found 212.143.

(4*R*)-4-Ethyl-8,8-diphenyl-6,10-dioxaspiro[4.5]decane-2-one (9). Purification by column chromatography (SiO₂ pentane/diethyl ether, 3:1, R_f = 0.24) gave 52 mg (31%) of **9** as a colorless liquid. ¹H-NMR (300 MHz) δ = 7.47-7.06 (m, 10H), 4.59 (d, *J* = 11.6 Hz, 2H), 4.26 (m, 2H), 3.07 (d, *J* = 18.0 Hz, 1H), 2.60-2.05 (m, 4H), 1.59 (m, 1H), 1.30 (m, 1H), 0.80 (t, *J* = 7.4 Hz, 3H); ¹³C-NMR (200 MHz) δ = 213.1, 143.7, 143.4, 128.7, 128.5, 128.1, 126.9, 126.4, 126.3, 104.7, 70.7, 68.8, 47.5, 44.8, 44.7, 42.8, 20.6, 12.4; HRMS calc. for C₂₂H₂₄O₃ 336.172, found 336.172; The ee was determined by HPLC on a chiral stationary phase (DAICEL CHIRALPAK AD, *i*PrOH:hexane 10:90, 1 ml/min, RT, T_r = 6.71 min, T_r = 13.77 min)

4-Ethyl-3-(2-hydroxy-8,8-dimethyl-6,10-dioxaspiro[4.5]dec-3-en-2-yl)-8,8-dimethyl-6,10-dioxaspiro[4.5]decan-2-one (8b). Purification by column chromatography (SiO₂ pentane/diethyl ether, 2:3, R_f = 0.42) gave 12 mg (12%) of **8b** as a colorless liquid. ¹H-NMR (500 MHz) δ = 6.12 (d, *J* = 5.8 Hz, 1H), 5.98 (d, *J* = 5.8 Hz, 1H), 3.49 (m, 8H), 2.96 (d, *J* = 14.2 Hz, 1H), 2.82 (d, *J* = 18.0 Hz, 1H), 2.59 (d, *J* = 18.0 Hz, 1H), 2.42 (q, *J* = 6.3 Hz, 1H), 2.38 (d, *J* = 6.0 Hz, 1H), 2.08 (d, *J* = 14.2 Hz, 1H), 1.91 (m, 1H), 1.26 (m, 1H), 1.16 (s, 3H), 1.02 (s, 3H), 0.98 (t, *J* = 7.8 Hz, 3H), 0.92 (s, 3H), 0.81 (s, 3H); ¹³C-NMR (200 MHz) δ = 213.8, 139.9, 132.1, 106.9, 103.5, 83.1, 72.7, 72.4, 72.0, 71.8, 60.9, 48.1, 46.0, 45.4, 30.0, 23.1, 22.8, 22.4, 22.1, 12.4; Ms (EI) for C₂₂H₃₄O₆: *m/z* = 395 (M⁺).

4-Ethyl-3-(2-hydroxy-8,8-dimethyl-6,10-dioxaspiro[4.5]dec-3-en-2-yl)-8,8-dimethyl-6,10-dioxaspiro[4.5]decan-2-one (8a). Purification by column chromatography (SiO₂ pentane/diethyl ether, 7:1, R_f = 0.38) gave 30 mg (30%) of **8a** as a colorless liquid. ¹H-NMR (500 MHz) δ = 6.30 (d, *J* = 5.8 Hz, 1H), 6.18 (d, *J* = 5.8 Hz, 1H), 3.52 (m, 8H), 2.92 (dd, *J* = 17.4 Hz, *J* = 0.9 Hz, 1H), 2.54 (d, *J* = 17.4 Hz, 1H), 2.47 (d, *J* = 8.1 Hz, 1H), 2.31 (q, *J* = 8.1 Hz, 1H), 2.27 (d, *J* = 14.0 Hz, 1H), 2.16 (d, *J* = 14.0 Hz, 1H), 1.91 (m, 1H), 1.36 (m, 1H), 1.15 (s, 3H), 1.00 (t, *J* = 7.8 Hz, 3H), 0.98 (s, 3H), 0.93 (s, 3H), 0.77 (s, 3H); ¹³C-NMR (200 MHz) δ = 213.8, 139.9, 132.1, 106.9, 103.5, 83.1, 72.7, 72.4, 72.0, 71.8, 60.9, 48.1, 46.0, 45.4, 30.0, 23.1, 22.8, 22.4, 22.1, 12.4; Ms (EI) for C₂₂H₃₄O₆: *m/z* = 395 (M⁺).

General procedure for the tandem 1,4-addition-aldol reaction (11a, 11b, 13a-21a).

A solution of Cu(OTf)₂ (3.6 mg, 0.01 mmol) and phosphoramidite¹ (0.02 mmol) in toluene (7 ml) was stirred under a nitrogen atmosphere at ambient temperature for 1 h. Cyclopenten-3,5-dione monoacetal (0.5 mmol) and the aldehyde (0.5 mmol) were added. After cooling the reaction mixture to -45 °C, the dialkylzinc reagent (0.6 ml, 1M solution in toluene) was added and stirring at -45°C was continued for 18 h. After complete conversion, the reaction mixture was poured in 25 ml NH₄Cl (aq), the organic layer was separated, and the aqueous layer was extracted two times with diethyl ether. The combined organic layers were dried over MgSO₄, filtered and concentrated *in vacuo*.

(3S,4R)-4-Ethyl-3-[(R)-hydroxy(phenyl)methyl]-8,8-dimethyl-6,10-dioxaspiro[4.5]decan-2-one (13a). Purification by column chromatography (SiO₂ hexane/diethyl ether, 1:2, R_f = 0.32) gave 107 mg (67%) of **13a** as a colorless oil which solidified upon standing. ¹H-NMR (300 MHz) δ = 7.28-7.23 (m, 5H), 4.75 (d, *J* = 8.1 Hz, 1H), 3.50 (s, 2H), 3.37 (d, *J* = 2.0 Hz, 2H), 2.74 (dd, *J* = 18.0 Hz, *J* = 1.2 Hz, 1H), 2.50 (d, *J* = 18 Hz, 1H), 2.46 (t, *J* = 7.5 Hz, 1H), 2.00 (q, *J* = 6.3 Hz, 1H), 1.55 (m, *J* = 6.3 Hz, 2H), 1.02 (s, 3H), 0.79 (s, 3H), 0.57 (t, *J* = 7.2 Hz, 3H); ¹³C-NMR (300 MHz) δ = 216.4, 140.8, 128.0, 127.7, 126.7, 103.2, 74.9, 71.7, 71.3, 59.1, 47.7, 44.7, 29.6, 22.2, 21.7, 20.6, 11.4; HRMS calc. for C₁₉H₂₆O₄ 318.183, found 318.183

(3S,4R)-4-Butyl-3-[(R)-hydroxy(phenyl)methyl]-8,8-dimethyl-6,10-dioxaspiro[4.5]decan-2-one (14a). Purification by column chromatography (SiO₂

(1) Arnold, L. A.; Imbos, R.; Mandoli, A.; de Vries, A. H. M.; Naasz, R.; Feringa, B. L. *Tetrahedron* **2000**, *56*, 2865.

hexane/diethyl ether, 1:1, R_f = 0.33) gave 111 mg (64%) of **14a** as a colorless oil which solidified upon standing. $^1\text{H-NMR}$ (300 MHz) δ = 7.31- 7.26 (m, 5H), 4.75 (d, J = 8.1 Hz, 1H), 3.97 (d, J = 1.5 Hz, 1H(OH)), 3.44 (s, 2H), 3.37 (d, J = 4.4 Hz, 2H), 2.86 (dd, J = 18.0 Hz, J = 1.2 Hz, 1H), 2.54 (d, J = 18.0 Hz, 1H), 2.40 (t, J = 7.5 Hz, 1H), 2.03 (q, J = 6.3 Hz, 1H), 1.52 (m, J = 6.6 Hz, 1H), 1.03 (s, 3H), 0.94 (m, 5H), 0.80 (s, 3H), 0.66 (m, 3H); $^{13}\text{C-NMR}$ (200 MHz) δ = 216.8, 141.0, 128.3, 128.1, 127.0, 103.5, 75.3, 72.1, 71.6, 59.9, 46.4, 45.1, 30.0, 29.5, 27.9, 22.6, 22.1, 21.6, 13.9; HRMS calc. for $\text{C}_{21}\text{H}_{30}\text{O}_4$ 346.214, found 346.213.

(3*S*,4*R*)-4-Ethyl-3-[(*R*)-hydroxy(phenyl)methyl]-8,8-diphenyl-6,10-dioxaspiro[4.5]decan-2-one (15a). Purification by column chromatography (SiO_2 pentane/diethyl ether, 2:1, R_f = 0.18) gave 167 mg (76%) of **15a** as a colorless oil which solidified upon standing. $^1\text{H-NMR}$ (300 MHz) δ = 7.31-7.02 (m, 15H), 4.74 (dd, J = 7.8 Hz, J = 1.8 Hz, 1H), 4.76-4.13 (m, 4H), 3.77 (d, J = 1.8 Hz, 1H(OH)), 2.52 (d, J = 18.0 Hz, 1H), 2.50 (d, J = 17.7 Hz, 1H), 2.40 (t, J = 7.2 Hz, 1H), 1.93 (q, J = 7.2 Hz, 1H), 1.37 (m, J = 7.2 Hz, 1H), 0.76 (m, J = 7.2 Hz, 1H), 0.31 (t, J = 7.2 Hz, 3H); $^{13}\text{C-NMR}$ (200 MHz) δ = 216.2, 205.8, 143.5, 143.3, 141.0, 128.6, 128.3, 128.3, 218.1, 127.9, 126.9, 126.7, 126.5, 125.8, 104.2, 75.2, 69.9, 69.1, 59.4, 48.0, 45.6, 44.9, 20.9, 11.7; Ms (CI) for $\text{C}_{29}\text{H}_{30}\text{O}_4$: m/z = 442 (M^+), 460 ($\text{M} + \text{NH}_4^+$).

(3*S*,4*R*)-3-[(*R*)-(4-Bromophenyl)(hydroxy)methyl]-4-ethyl-8,8-diphenyl-6,10-dioxaspiro[4.5]decan-2-one (17a). Purification by column chromatography (SiO_2 pentane/diethyl ether, 2:1, R_f = 0.31) gave 179 mg (69%) of **17a** as a colorless oil which solidified upon standing. $^1\text{H-NMR}$: (300 MHz) δ = 7.42-7.06 (m, 14H), 4.76 (d, J = 7.8 Hz, 1H), 4.51-4.17 (m, 4H), 2.96 (d, J = 18.0 Hz, 1H), 2.57 (d, J = 18.0 Hz, 1H), 2.40 (t,

$J = 7.2$ Hz, 1H), 1.94 (q, $J = 7.2$ Hz, 1H), 1.50 (m, $J = 7.2$ Hz, 1H), 0.85 (m, $J = 7.2$ Hz, 1H), 0.39 (t, $J = 7.2$ Hz, 3H); ^{13}C -NMR: (200 MHz) $\delta = 215.8, 143.4, 143.2, 140.1, 131.5, 128.6, 128.6, 128.3, 127.9, 126.9, 126.7, 126.5, 121.9, 104.1, 74.5, 69.9, 69.1, 59.2, 48.1, 45.4, 44.9, 20.9, 11.7$; HRMS calc. for $\text{C}_{29}\text{H}_{29}\text{O}_4\text{Br}$ 520.124, found 520.123.

(3*S*,4*R*)-3-[(*R*)-(4-Bromophenyl)(hydroxy)methyl]-4-butyl-8,8-diphenyl-6,10-dioxaspiro[4.5]decan-2-one (11a). Purification by column chromatography (SiO_2 pentane/diethyl ether, 2:1, $R_f = 0.28$) gave 169 mg (62%) of **11a** (major product) as colorless oil which solidified upon standing. ^1H -NMR: (300 MHz) $\delta = 7.42\text{--}7.05$ (m, 14H), 4.75 (dd, $J = 7.8$ Hz, $J = 1.8$ Hz, 1H), 4.53–4.46 (m, 2H), 4.31–4.16 (m, 2H), 3.84 (d, $J = 1.8$ Hz, 1H (OH)), 3.03 (dd, $J = 18.0$ Hz, $J = 1.8$ Hz, 1H), 2.53 (d, $J = 18.3$ Hz, 1H), 2.39 (dd, $J = 7.5$ Hz, $J = 7.2$ Hz, 1H), 2.02 (q, $J = 7.2$ Hz, 1H), 1.40 (m, 1H), 0.84–0.69 (m, 5H), 0.57 (t, $J = 7.2$ Hz, 3H); ^{13}C -NMR: (200 MHz) $\delta = 215.9, 143.3, 143.1, 140.1, 131.4, 128.7, 128.6, 128.3, 127.9, 126.9, 126.6, 126.4, 121.9, 103.9, 74.4, 70.1, 68.9, 59.6, 47.0, 45.2, 44.8, 29.6, 27.6, 22.7, 13.7$; Ms (EI) for $\text{C}_{31}\text{H}_{33}\text{BrO}_4$: $m/z = 550$ (M^+). (For NOESY-NMR see supporting material)

(3*S*,4*R*)-3-[(*S*)-(4-Bromophenyl)(hydroxy)methyl]-4-butyl-8,8-diphenyl-6,10-dioxaspiro[4.5]decan-2-one (11b). Purification by column chromatography (SiO_2 pentane/diethyl ether, 2:1, $R_f = 0.24$) gave 4 mg (1.5%) of **11b** (minor product) as colorless oil which solidified upon standing: ^1H -NMR: (300 MHz) $\delta = 7.42\text{--}7.00$ (m, 14H), 5.11 (dd, $J = 6.6$ Hz, $J = 3.6$ Hz, 1H), 4.55 (m, 2H), 4.28–4.12 (m, 2H), 3.49 (d, $J = 6.6$ Hz, 1H (OH)), 3.05 (dd, $J = 17.7$ Hz, $J = 1.8$ Hz, 1H), 2.50 (dd, $J = 9.3$ Hz, $J = 3.6$ Hz, 1H), 2.30 (d, $J = 17.7$ Hz, 1H), 2.15 (q, $J = 7.2$ Hz, 1H), 1.39 (m, 1H), 0.86–0.67 (m, 5H), 0.55 (t, $J = 7.2$ Hz, 3H); ^{13}C -NMR: (200 MHz) $\delta = 214.5, 143.2, 143.0, 140.9,$

131.4, 128.7, 128.3, 128.1, 127.6, 127.0, 126.4, 121.3, 103.9, 72.1, 70.5, 68.7, 60.0, 45.2 (two peaks), 44.7, 29.7, 27.5, 22.9, 13.7; Ms (EI) for C₃₁H₃₃BrO₄: m/z = 550 (M⁺). (For NOESY-NMR see supporting material)

(3*S*,4*R*)-4-Ethyl-3-[(1*S*)-1-hydroxybutyl]-8,8-diphenyl-6,10-dioxaspiro[4.5]decan-2-one (18a). Purification by column chromatography (SiO₂ pentane/diethyl ether, 3:2, R_f = 0.23) gave 133 mg (65%) of **18a** as a colorless oil which solidified upon standing; ¹H-NMR: (300 MHz) δ = 7.41-7.02 (m, 10H), 4.55 (m, 2H), 4.31-4.14 (m, 2H), 3.67 (q, *J* = 4.8 Hz, 1H), 3.04 (dd, *J* = 17.7 Hz, *J* = 1.2 Hz, 1H), 2.47 (d, *J* = 17.7 Hz, 1H), 2.41 (d, *J* = 5.1 Hz, 1H(OH)), 2.20 (m, 2H), 1.72- 1.20 (m, 7H), 0.88 (t, *J* = 7.2 Hz, 3H), 0.71 (t, *J* = 7.2 Hz, 3H); ¹³C-NMR: (200 MHz) δ = 214.9, 143.5, 143.2, 128.7, 128.3, 128.2, 126.9, 126.5, 126.3, 103.8, 71.4, 70.4, 68.8, 58.0, 49.4, 45.5, 44.7, 37.7, 20.8, 19.1, 13.9, 12.5; HRMS calc. for C₂₆H₃₂O₄ 408.230, found 408.232.

(8*S*,9*R*)-9-Ethyl-8-[(*R*)-hydroxy(phenyl)methyl]-2,2,3,3-tetramethyl-1,4-dioxaspiro[4.4]nonan-7-one (19a). Purification by column chromatography (SiO₂ pentane/diethyl ether, 2:1, R_f = 0.36) gave 90 mg (54%) of **19a** as a colorless oil. ¹H-NMR (300 MHz) δ = 7.35-7.22 (m, 5H), 4.85 (d, *J* = 9.6 Hz, 1H), 2.74 (d, *J* = 18.0 Hz, 1H), 2.52 (d, *J* = 18.0 Hz, 1H), 2.36 (d, *J* = 9.3 Hz, 1H), 1.68 (m, 1H), 1.20 (s, 3H), 1.19 (s, 3H), 1.17 (m, 1H), 1.16 (s, 3H), 1.12 (s, 3H), 0.92 (m, 1H), 0.40 (t, *J* = 7.2 Hz, 3H); ¹³C-NMR: (200 MHz) δ = 218.1, 141.1, 128.2, 128.0, 127.4, 110.4, 83.5, 83.3, 74.8, 60.4, 52.3, 50.5, 23.7, 23.5, 11.1; Ms (CI) for C₂₀H₂₈O₄: m/z = 332 (M⁺), 350 (M+NH₄⁺).

(8*S*,9*R*)-9-Ethyl-8-[(*R*)-hydroxy(phenyl)methyl]-1,4-dioxaspiro[4.4]nonan-7-one (20a). Crude product; ¹H-NMR (300 MHz) δ = 7.37-7.34 (m, 5H), 4.68 (d, *J* = 8.6 Hz, 1H), 3.95 (m, 4H), 2.68 (d, *J* = 13.6 Hz, 1 H), 2.50 (d, *J* = 13.6 Hz, 1 H), 2.49 (t, 1H),

1.92 (q, $J = 6.2$ Hz, 1H), 1.52 (m, $J = 6.5$ Hz, 1H), 0.93 (m, $J = 6.5$ Hz, 1H), 0.59 (t, $J = 7.2$ Hz, 3H); ^{13}C NMR (200 MHz) $\delta = 216.6, 140.9, 128.4, 128.1, 127.0, 112.9, 75.0, 65.0, 64.3, 60.5, 48.3, 48.1, 22.0, 11.7$; Ms (CI) for $\text{C}_{16}\text{H}_{20}\text{O}_4$: $m/z = 276$ (M^+), 294 ($\text{M}+\text{NH}_4^+$).

(2*S*,3*R*)-3-Ethyl-2-[(*R*)-hydroxy(phenyl)methyl]-4,4-dimethoxycyclopentanone

(21a). Crude product; ^1H -NMR (300 MHz) $\delta = 7.34\text{--}7.19$ (m, 5H), 4.72 (d, $J = 9.6$ Hz, 1H), 3.16 (d, $J = 9.9$ Hz, 6H), 2.61 (d, $J = 17.7$ Hz, 2H), 2.32 (m, 1H), 1.81, (m, 1H), 1.50 (m, 1H), 0.91 (m, 1H), 0.34 (t, $J = 7.2$ Hz, 3H); ^{13}C -NMR (200 MHz) $\delta = 217.1, 140.9, 128.2, 128.0, 127.3, 106.5, 75.5, 60.1, 50.1, 48.2, 46.0, 45.6, 23.2, 10.8$; Ms (CI) for $\text{C}_{16}\text{H}_{22}\text{O}_4$: $m/z = 278$ (M^+), 296 ($\text{M}+\text{NH}_4^+$).

General procedure for the oxidation to a diketone (11c, 13c-17c). To hydroxy ketone (0.2 mmol) in CH_2Cl_2 (5 ml) were added molecular sieves (4 Å, 0.5 g) and PCC (215 mg, 1 mmol) at 0°C . The reaction mixture was stirred for 4h at room temperature and diluted with diethyl ether, filtered over celite and evaporated to dryness.

(3*R*,4*R*)-3-Benzoyl-4-ethyl-8,8-dimethyl-6,10-dioxaspiro[4.5]decan-2-one (13c).

Purification by column chromatography (SiO_2 hexane/diethyl ether, 2;1, $R_f = 0.33$) gave 40 mg (64%) of **13c** as a colorless oil which solidified upon standing. ^1H -NMR (300 MHz) $\delta = 7.99$ (d, $J = 6.0$ Hz, 2H), 7.57-7.43 (m, 3H), 4.26 (d, $J = 11.4$ Hz, 1H), 3.58-3.44 (m, 4H), 3.25 (d, $J = 17.7$ Hz, 1H), 3.14 (m, 1H), 2.43 (d, $J = 17.4$ Hz, 1H), 1.97 (m, 1H), 1.54 (m, 1H), 1.22 (s, 3H), 0.77 (t, $J = 7.5$ Hz, 3H), 0.75 (s, 3H); ^{13}C -NMR (200 MHz) $\delta = 207.1, 195.2, 137.1, 133.5, 129.5, 128.7, 102.5, 73.5, 71.2, 62.3, 50.9, 44.6, 30.1, 22.6, 21.9, 20.1, 12.9$; HRMS calc. for $\text{C}_{19}\text{H}_{24}\text{O}_4$ 316.168, found 316.167; the ee of

87 % was determined by HPLC on a chiral stationary phase (DAICEL CHIRALPAK AD, *i*PrOH:hexane 3:97, 1 ml/min, RT, T_r = 17.0 min, T_r = 20.9 min).

(3*R*,4*R*)-3-Benzoyl-4-butyl-8,8-dimethyl-6,10-dioxaspiro[4.5]decan-2-one (14c).

Purification by column chromatography (SiO₂ hexane/diethyl ether, 5:2, R_f = 0.38) gave 43 mg (63%) of **14c** as a colorless oil which solidified upon standing. ¹H-NMR (300 MHz) δ = 7.98 (d, J = 7.5 Hz, 2H), 7.50 (m, 3H), 4.26 (d, J = 11.0 Hz, 1H), 3.45 (m, 4H), 3.25 (d, J = 17.0 Hz, 1H), 2.43 (d, J = 17.0 Hz, 1H), 1.92 (m, 1H), 1.46 (m, 1H), 1.22 (m, 7H), 1.04 (m, 1H), 0.77 (t, J = 7 Hz, 3H), 0.75 (s, 3H); ¹³C-NMR (200 MHz) δ = 207.0, 194.9, 137.1, 133.4, 129.4, 128.6, 102.5, 73.5, 71.2, 62.7, 49.2, 44.5, 30.5, 30.2, 26.9, 22.9, 22.8, 22.0, 13.9; HRMS calc. for C₂₁H₂₈O₄ 344.198, found 344.199; the ee of 87 % was determined by HPLC on a chiral stationary phase (DAICEL CHIRALPAK AD, *i*PrOH:hexane 10:90, 1ml/min, RT, T_r = 7.8 min, T_r = 11.1)

(3*R*,4*R*)-3-Benzoyl-4-ethyl-8,8-diphenyl-6,10-dioxaspiro[4.5]decan-2-one (15c).

Purification by column chromatography (SiO₂ pentane/diethyl ether, 5:1, R_f = 0.30) gave 67 mg (76%) of **15c** as a colorless oil which solidified upon standing. ¹H-NMR (300 MHz) δ = 7.99 (d, J = 6.0 Hz, 2H), 7.57-7.03 (m, 13H), 4.65 (d, J = 11.7 Hz, 2H), 4.35-4.21 (m, 3H), 3.37 (d, J = 17.7 Hz, 1H), 3.11 (m, 1H), 2.56 (d, J = 17.1 Hz, 1H), 1.59 (m, 1H), 1.30 (m, 1H), 0.57 (t, J = 6.9 Hz, 3H); ¹³C-NMR (200 MHz) δ = 206.6, 194.8, 143.4, 143.1, 136.9, 133.5, 129.4, 128.7, 128.6, 128.1, 127.0, 126.3, 103.0, 71.4, 68.6, 62.4, 50.8, 45.1, 44.7, 20.1, 12.8; enol form <1%; HRMS calc. for C₂₉H₂₈O₄ 440.198, found 440.197. The ee of 94 % was determined by HPLC on a chiral stationary phase (DAICEL CHIRALPAK AD, *i*PrOH:hexane 50:50, 1 ml/min, RT, T_r = 5.64 min, T_r = 13.11 min).

(3*R*,4*R*)-3-(4-Bromobenzoyl)-4-ethyl-8,8-diphenyl-6,10-dioxaspiro[4.5]decan-2-one (17c). Purification by column chromatography (SiO₂ pentane/diethyl ether, 5:1, R_f = 0.36) gave 77 mg (76%) of **17c** as a colorless oil which solidified upon standing. ¹H-NMR (300 MHz) δ = 7.83 (d, *J* = 8.4 Hz, 2H), 7.63-7.03 (m, 12H), 4.65 (d, *J* = 12.0 Hz, 2H), 4.44-4.16 (m, 4H), 3.36 (d, *J* = 17.4 Hz, 1H), 3.11 (m, 1H), 2.54 (d, *J* = 17.1 Hz, 1H), 1.59 (m, 1H), 1.31 (m, 1H), 0.55 (t, *J* = 6.9 Hz, 3H); ¹³C-NMR (200 MHz) δ = 206.3, 193.7, 143.4, 143.0, 135.6, 131.9, 130.9, 129.0, 128.8, 128.6, 128.1, 127.1, 126.4, 126.3, 102.9, 71.4, 68.6, 62.4, 50.6, 45.0, 44.7, 20.1, 12.8; HRMS calc. for C₂₉H₂₇O₄ Br 518.109, found 518.107. The ee of 96 % was determined by HPLC on a chiral stationary phase (DAICEL CHIRALPAK AD, *i*PrOH:hexane 50:50, 1 ml/min, RT, T_r = 6.7 min, T_r = 11.9 min).

(3*R*,4*R*)-3-(4-Bromobenzoyl)-4-butyl-8,8-diphenyl-6,10-dioxaspiro[4.5]decan-2-one (11c). Purification by column chromatography (SiO₂ pentane/diethyl ether, 5:1, R_f = 0.30) gave 68 mg (62%) of **11c** as a colorless oil which solidified upon standing. ¹H-NMR (300 MHz) δ = 7.82-7.02 (m, 14H), 4.62 (d, *J* = 11.7 Hz, 2H), 4.41-4.13 (m, 2H), 3.33 (d, *J* = 17.7 Hz, 1H), 3.10 (m, 1H), 2.48 (d, *J* = 17.7 Hz, 1H), 1.51 (m, 1H), 1.22-0.74 (m, 5H), 0.62 (t, *J* = 6.9 Hz, 3H); ¹³C-NMR (200 MHz) δ = 206.3, 193.5, 143.2, 142.9, 135.6, 131.9, 130.9, 128.7, 128.6, 128.1, 127.1, 126.3, 126.2, 102.9, 71.4, 68.6, 62.7, 48.8, 44.9, 44.7, 30.2, 27.0, 22.8, 13.6; HRMS calc. for C₃₁H₃₁O₄Br 548.138 found 548.139. The ee of 97 % was determined by HPLC on a chiral stationary phase (DAICEL CHIRALPAK AD, *i*PrOH:hexane 50:50, 1 ml/min, RT, T_r = 5.70 min, T_r = 9.43 min).

(4*R*,5*S*)-4-Ethyl-3-(3-hydroxy-2,2-dimethylpropoxy)-5-[(*R*)-hydroxy(phenyl)methyl]-2-cyclopenten-1-one (13d). Purification by column

chromatography (SiO₂ MeOH/diethyl ether, 1:40, R_f = 0.37) gave 107 mg (67%) of **13d** as a white solid. ¹H-NMR (300 MHz) δ = 7.36-7.21 (m, 5H), 5.27 (s, 1H), 5.20 (s, 1H(OH)), 4.58 (d, *J* = 9.6 Hz, 1H), 3.78 (d, *J* = 9.2 Hz, 1H), 3.67 (d, *J* = 9.2 Hz, 1H), 3.41 (m, 2H), 2.39 (m, 2H), 1.51 (s, 3H), 1.21 (m, 1H), 0.93 (s, 3H), 0.87 (m, 1H), 0.46 (t, *J* = 7.5 Hz, 3H); HRMS calc. for C₁₉H₂₆O₄ 318.183, found 318.181.

(4*R*,5*S*)-4-Ethyl-3-(2-hydroxyethoxy)-5-[(*R*)-hydroxy(phenyl)methyl]-2-cyclopenten-1-one (20d). Purification by column chromatography (SiO₂ MeOH/diethyl ether, 1:30, R_f = 0.31) gave 104 mg (75%) of **20d** as a white solid. ¹H-NMR (300 MHz) δ = 7.35-7.21 (m, 5H), 5.30 (s, 1H), 5.10 (s, 1H(OH)), 4.59 (d, *J* = 9.9 Hz, 1H), 4.04 (m, 2H), 3.91 (m, 2H), 2.41 (m, 2H), 1.20 (m, *J* = 7.2 Hz, 1H), 0.96 (m, 1H), 0.44 (t, *J* = 7.5 Hz, 3H); ¹³C-NMR (200 MHz) δ = 208.2, 192.4, 141.2, 128.4, 128.3, 127.0, 103.7, 76.4, 73.3, 60.3, 55.7, 44.0, 22.8, 8.9; HRMS calc. for C₁₆H₂₀O₄ 276.135, found 276.136.

expi relayh

SAMPLE

DEC. 6 VT

date Mar 16 2001 dirq 499.862

solvent CDC13 dn

file /dsk3/nmr/org-dprw H1

/legct/PDE1-methyl- dof 38

ester-COBSY-500pfg- dm 0

sw-160301.f1d dnm nmh

ACQUISITION c 9000

sfreq 499.861 dseq

tn H1 dres 1.0

at 0.228 homo n

np 2048 PROCESSING

sw 4482.8 sb 0.114

fb 2600 sbs not used

ss 2 wfilie

cpwr 60 proc ft

pw 16.0 fn 2048

pl 16.0 math t

d1 1.000

tof -689.6 weir

nt 4 wexp

ct 4 wbs

tau 0 wnt

relay 0 2D PROCESSING

alock n sb1 0.029

gain 26 sbs1 not used

FLAGS wfilie1

l1 n procl ft

ln n fn1 2048

dp Y

hs nm

2D ACQUISITION

sw1 4482.8

nl 256

phase 1

DISPLAY

sp 374.5

wp 2524.0

vs 583



sc 6


wc 100

hzmm 34.48


ls 815.76


Prostaglandin E1 methyl ester (28)







Prostaglandin E1 methyl ester (28)







4,4-Dimethoxy-2-cyclopenten-1-one (**7a**)










1,4-Dioxaspiro[4.4]non-8-en-7-one (**7c**)






INDEX	FREQUENCY	PPM	HEIGHT
1	15332.909	203.267	14.7
2	11803.641	156.480	41.3
3	10819.694	143.436	18.6
4	10803.824	143.225	23.3
5	10193.435	135.133	40.4
6	9684.371	128.385	120.9
7	9669.722	128.190	107.6
8	9623.332	127.576	118.5
9	9557.410	126.702	126.0
10	9554.969	126.669	66.9
11	9534.216	126.394	50.1
12	7848.322	104.044	24.2
13	5840.142	77.422	5.6
14	5808.402	77.001	5.3
15	5775.441	76.564	5.5
16	5298.117	70.237	89.1
17	3348.535	44.391	22.4
18	3319.237	44.003	39.0






2,2,3,3-Tetramethyl-1,4-dioxaspiro[4.4]non-8-en-7-one


(7e)







(4R)-4-Ethyl-8,8-dimethyl-6,10-dioxaspiro[4.5]decane-2-one

(8)





4-Ethyl-3-(2-hydroxy-8,8-dimethyl-6,10-dioxaspiro[4.5]dec-3-en-2-yl)-8,8-dimethyl-6,10-dioxaspiro[4.5]decan-2-one

(8a)




4-Ethyl-3-(2-hydroxy-8,8-dimethyl-6,10-dioxaspiro[4.5]dec-3-en-2-yl)-8,8-dimethyl-6,10-dioxaspiro[4.5]decan-2-one

(8b)





INDEX FREQUENCY PPM HEIGHT
 /export/home/10718.080 213.147
 2 7227.144 143.724 9.9

3	7208.833	143.360	18.9
4	6470.285	128.672	106.3
5	6459.985	128.467	109.2
6	6442.437	128.118	107.5
7	6382.926	126.935	63.7
8	6358.511	126.449	103.5
9	6351.263	126.305	61.7
10	5262.897	104.661	36.2
11	3903.679	77.631	53.9
12	3872.016	77.001	53.4
13	3839.971	76.364	54.3
14	3555.386	70.705	51.9
15	3460.779	68.823	51.6
16	2389.198	47.513	62.9
17	2252.627	44.797	42.4
18	2249.957	44.744	63.6
19	2153.824	42.832	64.7
20	1035.320	20.589	65.0
21	622.939	12.388	62.6





(4R)-4-Ethyl-8,8-diphenyl-6,10-dioxaspiro[4.5]decane-2-one (9)







(4R)-4-Butyl-8,8-diphenyl-6,10-dioxaspiro[4.5]decan-2-one (10)





3-[(4-bromophenyl)(hydroxymethyl)-4-butyl-8,8-diphenyl-6,10-dioxaspiro[4.5]decan-2-one (IIa)







(3S,4R)-4-Ethyl-3-[(R)-hydroxy(phenyl)methyl]-8,8-dimethyl-6,10-dioxaspiro[4.5]decan-2-one

(13a)


PULSE SEQUENCE
 Relax. delay 1.000 sec
 Pulse 90.0 degrees
 Acq. time 1.498 sec
 Width 12500.0 Hz
 1024 repetitions
 OBSERVE C13, 50.2811248 MHz
 DECOUPLE H1, 199.9652953 MHz
 Power 39 dB
 continuously on
 WALTZ-16 modulated
 DATA PROCESSING
 Line broadening 1.5 Hz
 FT size 65536
 Total time 42 minutes






(3S,4R)-4-butyl-3-[(R)-hydroxy(phenyl)methyl]-8,8-dimethyl-6,10-dioxaspiro[4.5]decan-2-one

(14a)




INDEX FREQUENCY PPM HEIGHT
1 10869.147 216.151 7.9
2 10348.044 205.788 11.3
3 10348.044 205.788 11.3
4 7205.019 143.284 11.3
5 7089.430 140.985 16.8
6 6467.234 128.612 -79.0
7 6453.882 128.346 -94.5
8 6450.448 128.278 -88.6
9 6439.385 128.058 -88.2
10 6431.756 127.906 -80.5
11 6382.163 126.920 -92.9
12 6372.626 126.730 -81.9
13 6358.893 126.457 -47.1
14 6325.704 125.797 -5.7
15 5237.338 104.153 20.2
16 3904.060 77.639 24.8
17 3872.016 77.001 25.3
18 3840.353 76.372 24.3
19 3778.934 75.150 -39.9
20 3516.475 69.931 34.2
21 3471.842 69.043 35.0
22 2989.268 59.447 -44.6
23 2412.468 47.976 -39.9
24 2293.064 45.601 35.4
25 2255.679 44.858 23.4
26 1050.580 20.893 36.8
27 587.080 11.675 -33.4




(3S,4R)-4-Ethyl-3-[(1R)-hydroxy(phenyl)methyl]-8,8-diphenyl-6,10-dioxaspiro[4.5]decan-2-one

(15a)






(3S,4R)-4-Butyl-3-[(R)-hydroxy(phenyl)methyl]-8,8-diphenyl-6,10-dioxaspiro[4.5]decan-2-one


(16a)



INDEX	FREQUENCY	PPM	HEIGHT
1	10877.548	216.318	18.3
2	10865.332	216.075	16.7
3	7209.596	143.375	22.3
4	7202.730	143.238	-4.3
5	7087.522	140.947	73.2
6	6529.796	129.856	88.1
7	6468.378	128.634	92.7
8	6455.408	128.376	49.1
9	6450.448	128.278	83.3
10	6440.911	128.088	126.0
11	6432.900	127.929	81.1
12	6381.019	126.897	48.6
13	6368.430	126.647	17.9
14	6356.985	126.419	51.6
15	5231.615	104.039	49.9
16	3903.679	77.631	50.8
17	3871.634	76.994	38.3
18	3839.971	76.364	32.4
19	3781.605	75.203	42.7
20	3525.249	70.105	33.7
21	3463.449	68.876	43.0
22	3006.816	59.795	36.3
23	2356.009	46.853	17.0
24	2284.290	45.427	48.7
25	2254.916	44.843	40.4
26	1487.757	29.587	48.6
27	1390.861	27.660	40.5
28	1140.609	22.683	
29	692.750	13.776	









(3S,4R)-4-ethyl-3-[(1S)-1-hydroxybutyl]-8,8-diphenyl-6,10-dioxaspiro[4.5]decan-2-one


(18a)






(8S,9R)-9-Ethyl-8-[(R)-hydroxy(phenyl)methyl]-2,2,3,3-tetramethyl-1,4-dioxaspiro[4.4]nonan-7-one


(19a)







(8S,9R)-9-Ethyl-8-[(R)-hydroxy(phenyl)methyl]-1,4-dioxaspiro[4.4]nonan-7-one


(20a)






(2S,3R)-3-Ethyl-2-[(R)-hydroxy(phenyl)methyl]-4,4-dimethoxycyclopentanone
(21a)





3-(4-bromobenzoyl)-4-butyl-8,8-diphenyl-6,10-dioxaspiro[4.5]decan-2-one (IIc)



/export/home/vmactl/vmactls/data/auto/ch/0202

Solvent: CDCl₃

Ambient temperature

Sample #2

File: 202

GENI-200NB "smc200"

PULSE SEQUENCE

Relax. delay 1.000 sec

Pulse 90.0 degrees

Acq. time 1.498 sec

Width 12500.0 Hz

2200 repetitions

OBSERVE C13, 50.281248 MHz

DECOUPLE H1, 199.9652953 MHz

Power 39 dB

continuously on


WALTZ-16 modulated

DATA PROCESSING

Line broadening 1.5 Hz


FT size 65536


Total time 91 minutes



(13c)


(3R,4R)-3-Benzoyl-4-ethyl-8,8-dimethyl-6,10-dioxaspiro[4.5]decan-2-one






(3R,4R)-3-Benzoyl-4-butyl-8,8-dimethyl-6,10-dioxaspiro[4.5]decan-2-one

(14c)






(3R,4R)-3-Benzoyl-4-ethyl-8,8-diphenyl-6,10-dioxaspiro[4.5]decan-


2-one

(15c)




3R,4R)-3-(4-Bromobenzoyl)-4-ethyl-8,8-diphenyl-6,10-dioxaspiro[4.5]decan-2-one


(17c)



(4R,5S)-4-Ethyl-3-(3-hydroxy-2,2-dimethylpropoxy)-5-(1(R)-hydroxy(phenyl)methyl)-2-cyclopenten-1-one



(13d)



Export/home/vnmr1/vnmrsys/data/auto/tu/0201

Solvent: CDCl₃

Ambient temperature

Sample #2

File: 201

GEMINI-200B "nmr200"

PULSE SEQUENCE

Relax. delay 1.000 sec

Pulse 90.0 degrees

Acq. time 1.498 sec

Width 12500.0 Hz

1600 repetitions

OBSERVE C13, 50.281921 MHz

DECOUPLE H1, 199.9655124 MHz

Power 39 dB

continuously on


WALTZ-16 modulated

DATA PROCESSING

Line broadening 1.5 Hz


FT size 65536

Total time 65 minutes




(4R,5S)-4-Ethyl-3-(2-hydroxyethoxy)-5-[(R)-hydroxy(phenyl)methyl]-2-cyclopenten-1-one

(20d)





1A




(4R,5S)-4-Ethyl-5-[(R)-hydroxy(phenyl)methyl]-3-methoxy-2-cyclopenten-1-one


(21d)









(4S,5R)-5-ethyl-4-[(R)-hydroxy(phenyl)methyl]-2-cyclopenten-1-one
(23)







Bis(methyl-6-hepanoate)zinc
(24)










Prostaglandin E1 methyl ester (28)






(Z)-3-Iodo-2-octen-1-ol (**30**)






(Z)-3-dimethyl(phenyl)silyl-2-octen-1-ol (**32**)








(26)

Methyl 7-(((1R,2S)-2-(((1R,2Z)-3-
[dimethyl(phenyl)silyl]-1-hydroxy-2-octenyl]-3-oxo-
8,8-diphenyl-6,10-dioxaspiro[4.5]dec-1-yl))heptanoate








Methyl 7-(2-((Z)-3-[dimethyl(phenyl)silyl]-1-hydroxy-2-octenyl)-3-hydroxy-8,8-diphenyl-6,10-dioxaspiro[4.5]dec-1-yl)heptanoate (**35**)



(36)



Methyl 7-((1*R*,2*R*,3*R*)-3-(acetyloxy)-2-((*E*,3*S*)-3-(acetyloxy)-1-octenyl)-8,8-diphenyl-6,10-dioxaspiro[4.5]dec-1-yl)heptanoate
(37)



(38)

Methyl 7-((1R,2R,3R)-3-hydroxy-2-((E,3S)-3-hydroxy-1-octenyl)-8,8-diphenyl-6,10-dioxaspiro[4.5]dec-1-yl)heptanoate